



# The Breastfeeding-Friendly Pediatric Office Practice

Joan Younger Meek, MD, MS, RD, FAAP, IBCLC, Amy J. Hatcher, MD, FAAP, SECTION ON BREASTFEEDING

The landscape of breastfeeding has changed over the past several decades as more women initiate breastfeeding in the postpartum period and more hospitals are designated as Baby-Friendly Hospitals by following the evidence-based Ten Steps to Successful Breastfeeding. The number of births in such facilities has increased more than sixfold over the past decade. With more women breastfeeding and stays in the maternity facilities lasting only a few days, the vast majority of continued breastfeeding support occurs in the community. Pediatric care providers evaluate breastfeeding infants and their mothers in the office setting frequently during the first year of life. The office setting should be conducive to providing ongoing breastfeeding support. Likewise, the office practice should avoid creating barriers for breastfeeding mothers and families or unduly promoting infant formula. This clinical report aims to review practices shown to support breastfeeding that can be implemented in the outpatient setting, with the ultimate goal of increasing the duration of exclusive breastfeeding and the continuation of any breastfeeding.

## BACKGROUND/CURRENT RECOMMENDATIONS

Breastfeeding has long been documented as the ideal method for feeding and promoting the optimal development of infants and children, with rare exceptions (see Recommendation 3 below). The American Academy of Pediatrics (AAP) describes breastfeeding as the normative method of infant feeding. There are countless medical, emotional, and economic benefits of breastfeeding as described in the 2012 AAP statement "Breastfeeding and the Use of Human Milk."<sup>1</sup> Benefits of breastfeeding include decreased risk of lower respiratory infections, gastroenteritis, otitis media, and necrotizing enterocolitis, the latter being especially important in preterm infants. Because breastfeeding is the norm for infant feeding, comparatively there are risks associated with the lack of breastfeeding, which include an increase in sudden infant death syndrome, obesity, asthma, certain childhood cancers, diabetes, and postneonatal death.<sup>1-3</sup> Breastfeeding promotes attachment and optimal

## abstract

FREE

*Department of Clinical Sciences, Florida State University College of Medicine, Tallahassee, Florida*

*Drs Meek and Hatcher conceptualized the review, developed the search criteria, contributed to drafting the initial manuscript and to manuscript revisions, and responded to questions and comments from reviewers and the Board of Directors; and both authors approved the final manuscript as submitted.*

*This document is copyrighted and is property of the American Academy of Pediatrics and its Board of Directors. All authors have filed conflict of interest statements with the American Academy of Pediatrics. Any conflicts have been resolved through a process approved by the Board of Directors. The American Academy of Pediatrics has neither solicited nor accepted any commercial involvement in the development of the content of this publication.*

*Clinical reports from the American Academy of Pediatrics benefit from expertise and resources of liaisons and internal (AAP) and external reviewers. However, clinical reports from the American Academy of Pediatrics may not reflect the views of the liaisons or the organizations or government agencies that they represent.*

*The guidance in this statement does not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate.*

*All clinical reports from the American Academy of Pediatrics automatically expire 5 years after publication unless reaffirmed, revised, or retired at or before that time.*

**DOI:** 10.1542/peds.2017-0647

Address correspondence to Joan Younger Meek, MD, MS, RD, FAAP, IBCLC. E-mail: joan.meek@med.fsu.edu.

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2017 by the American Academy of Pediatrics

**FINANCIAL DISCLOSURE:** The authors have indicated they have no financial relationships relevant to this article to disclose.

**To cite:** Meek JY, Hatcher AJ, AAP SECTION ON BREASTFEEDING. The Breastfeeding-Friendly Pediatric Office Practice. *Pediatrics*. 2017;139(5):e20170647

cognitive development. In women, lack of breastfeeding is associated with an increase in the risk of breast and ovarian cancer, type 2 diabetes, heart disease, and postpartum depression.<sup>4-7</sup>

The AAP recommends exclusive breastfeeding for approximately 6 months, followed by continued breastfeeding for 1 year or longer, as mutually desired by mother and child.<sup>1</sup> The *Surgeon General's Call to Action To Support Breastfeeding*<sup>8</sup> in 2011 emphasized the importance of breastfeeding as a public health imperative. Breastfeeding is strongly promoted, supported, and encouraged by the AAP, the Academy of Breastfeeding Medicine,<sup>9</sup> the American College of Obstetricians and Gynecologists,<sup>10,11</sup> and the American Academy of Family Physicians.<sup>12</sup> Each of these organizations calls on its members to be actively engaged in promoting and supporting breastfeeding among their patients.

## EPIDEMIOLOGY

The rate of initiation of any breastfeeding in the US population is 81.1% according to data from the National Immunization Survey (2016, birth cohort from 2013).<sup>13</sup> Although the rate of breastfeeding initiation approaches the Healthy People 2020 target<sup>14</sup> of 81.9%, only 22.3% of US infants are breastfed exclusively at age 6 months. There are significant disparities in terms of breastfeeding rates in the country; among black infants, only 66.3% are breastfed at all, and only 14.6% are exclusively breastfed through the first 6 months of life. Among Native American and Alaska Native infants, breastfeeding initiation is 68.3% and exclusive breastfeeding rates at 6 months are 17.9%. Mothers are more likely to breastfeed if they are married, have a college education, live in metropolitan areas, do not experience poverty, and do not

**TABLE 1** The Ten Steps to Successful Breastfeeding

1. Have a written breastfeeding-friendly policy that is routinely communicated to all health care staff.
2. Train all health care staff in the skills necessary to implement this policy.
3. Inform all pregnant women about the benefits and management of breastfeeding.
4. Help mothers initiate breastfeeding within 1 hour of birth.
5. Show mothers how to breastfeed and how to maintain lactation, even if they are separated from their infants.
6. Give infants no food or drink other than breast milk, unless medically indicated.
7. Practice rooming in: allow mothers and infants to remain together 24 hours a day.
8. Encourage breastfeeding on demand.
9. Give no pacifiers or artificial nipples to breastfeeding infants.<sup>a</sup>
10. Foster the establishment of breastfeeding support groups and refer mothers to them on discharge from the hospital or birth center.

<sup>a</sup> The AAP does not support a categorical ban on pacifiers because of their role in SIDS risk reduction and their analgesic benefit during painful procedures when breastfeeding cannot provide the analgesia. Pacifier use in the hospital in the neonatal period should be limited to specific medical indications, such as pain reduction, calming in a drug-exposed infant, etc. For breastfed infants, pacifier introduction should be delayed until breastfeeding is firmly established. Infants who are not being directly breastfed can begin pacifier use as soon as desired.

receive benefits from the Special Supplemental Nutrition Program for Women, Infants, and Children (WIC). WIC provides targeted breastfeeding support and peer-counseling services for mothers who qualify for their services. Infants most likely to experience toxic stress are least likely to be breastfed. According to the 2005–2007 Infant Feeding Practices Study II, 85% of US mothers intended to breastfeed exclusively for  $\geq 3$  months; however, only 32.4% achieved their intended exclusive breastfeeding duration.<sup>15</sup>

## CURRENT INITIATIVES TO INCREASE BREASTFEEDING RATES

There have been numerous initiatives to increase breastfeeding exclusivity and duration both in hospitals and in the outpatient setting. US hospitals accredited by The Joint Commission with maternity units that have at least 1100 births per year are required to report data on the Perinatal Care Core Measure Set, which includes a measure on Exclusive Breast Milk Feeding.<sup>16</sup> There have been major initiatives to increase the number of maternity facilities designated as Baby-Friendly. The first of these was Best Fed Beginnings,<sup>17</sup> a collaboration of the National Initiative for Children's Healthcare Quality, the Centers for Disease Control and Prevention

(CDC), and Baby-Friendly USA, in a national effort to improve maternity care practices and achieve designation as Baby-Friendly. Through Best Fed Beginnings, hospitals implemented the evidence-based and AAP-endorsed Ten Steps to Successful Breastfeeding (Table 1), as established in the Baby-Friendly Hospital Initiative, developed by the World Health Organization (WHO)/United Nations Children's Fund.<sup>18-20</sup> Additional maternity care facilities are pursuing the Baby-Friendly designation by participating in EMPower Breastfeeding: Enhancing Maternity Practices,<sup>21</sup> in cooperation with the CDC's Division of Nutrition, Physical Activity, and Obesity, and in partnership with the Carolina Global Breastfeeding Institute and Population Health Improvement Partners. EMPower similarly provides support to hospitals in quality-improvement methodology and breastfeeding support to improve maternity care practices and achieve Baby-Friendly USA designation. The number of births in designated facilities increased from 2.9% in 2007 to 20.01% in January 2017,<sup>22</sup> surpassing the Healthy People 2020 goal of 8.1%.<sup>14</sup>

For the national breastfeeding targets to be met, outpatient support from pediatricians and other pediatric care providers is imperative. As more infants are

being discharged from hospitals designated as Baby-Friendly and having successfully initiated breastfeeding, it is essential that the pediatricians with whom they will follow up are knowledgeable about breastfeeding and that their office practices are prepared to support these breastfeeding dyads. The *Surgeon General's Call to Action To Support Breastfeeding* lists specific action steps that apply directly to the pediatric outpatient practice (Table 2).

Many pediatricians have little experience in clinical breastfeeding management. Analysis of an AAP Periodic Survey of Fellows regarding breastfeeding in 2004, compared with a similar survey in 1995, showed that pediatricians were less likely to believe that the benefits of breastfeeding outweighed the difficulties or inconvenience, and fewer believed that almost all mothers were able to succeed.<sup>23</sup> Unfortunately, more pediatricians in 2004 reported reasons to recommend against breastfeeding compared with the cohort who responded in 1995. A 2014 AAP Periodic Survey confirmed that some of these attitudes persist (unpublished data from American Academy of Pediatrics Periodic Survey of Fellows No. 89).

Szucs et al<sup>24</sup> found that there were gaps in providers' breastfeeding knowledge, counseling skills, and professional education and training. The authors showed that providers' cultures and attitudes affected breastfeeding promotion and support and that the providers used their own breastfeeding experiences to replace evidence-based knowledge and AAP policy statement recommendations. Both the Academy of Breastfeeding Medicine (ABM)<sup>25</sup> and the US Breastfeeding Committee<sup>26</sup> have published recommendations regarding the education and training of health care professionals, including pediatric care providers,

**TABLE 2** Relevant Steps in the US Surgeon General's Call to Action to Support Breastfeeding

1. Give mothers the support they need to breastfeed their infants.
6. Ensure that the marketing of infant formula is conducted in a way that minimizes its negative impacts on exclusive breastfeeding.
8. Develop systems to guarantee continuity of skilled support for lactation between hospitals and health care settings in the community.
9. Provide education and training in breastfeeding for all health professionals who care for women and children.
10. Include basic support for breastfeeding as a standard of care for midwives, obstetricians, family physicians, nurse practitioners, and pediatricians.
11. Ensure access to services provided by IBCLCs.

Shown are action steps that apply directly to the pediatric outpatient practice, numbered according to the original steps in the Call to Action.<sup>8</sup>

in breastfeeding support and management.

The ABM's clinical protocol "The Breastfeeding-Friendly Physician's Office: Optimizing Care for Infant and Children"<sup>27</sup> details specific steps that a practice can take to become more breastfeeding friendly. These guidelines can inform the guidance for pediatric offices. Corriveau et al<sup>28</sup> sought to determine whether implementing a program based on this clinical protocol affected breastfeeding rates within the pediatric primary care setting. Even with a diverse patient population, rates of both initiation of and exclusive breastfeeding increased after the implementation of the ABM breastfeeding-friendly protocol. Feldman-Winter<sup>29</sup> outlined methods to increase breastfeeding initiation and duration in "Evidence-Based Interventions To Support Breastfeeding." In a comprehensive evidence review for the US Preventive Services Task Force, Chung et al<sup>30</sup> provided evidence that breastfeeding interventions, including both pre- and postnatal interventions, were more influential than either alone and that interventions including a component of lay support, such as peer counselors, are more effective than usual care in increasing short- and long-term breastfeeding rates. The US Preventive Services Task Force recommends interventions during pregnancy and after birth to support breastfeeding and concludes that coordinated interventions can

increase breastfeeding initiation, duration, and exclusivity (Evidence Grade B).<sup>31</sup>

### THE BREASTFEEDING-FRIENDLY PEDIATRIC OFFICE PRACTICE

Because of the importance of breastfeeding to maternal and infant health outcomes, all pediatric care providers should aim to improve breastfeeding rates in their practices, with the goal of achieving the AAP breastfeeding recommendations for exclusive breastfeeding until approximately 6 months, followed by continued breastfeeding for 1 year or longer, as mutually desired by the mother and infant.<sup>1</sup> Pediatric care providers should help mothers identify and reach their own breastfeeding goals.

The following evidence-based recommendations for the pediatric outpatient practice should be considered as part of the practice-improvement process to increase breastfeeding rates to meet or exceed the AAP recommendations and the Healthy People 2020 goals. These recommendations provide guidance for clinical care and are not intended to imply a standard of care, nor does this report provide a strict weighing of the evidence:

1. Establish a written breastfeeding-friendly office policy that includes the provisions as outlined in this document and in Table 3.<sup>27,32,33</sup> Provide a lactation room with

supplies for employees who breastfeed or express breast milk at work. This room could also be used by breastfeeding mothers (refer to recommendation 12). Collaborate with the entire team, including colleagues and office staff. All employees should be aware of the policy, and copies of the policy should be provided to all staff, including anyone who answers the telephone or retrieves messages, such as front office staff.

2. Train staff in the skills necessary to support breastfeeding, especially nurses and medical assistants.<sup>33</sup> Identify  $\geq 1$  breastfeeding resource personnel on staff. If possible, consider employing an International Board Certified Lactation Consultant (IBCLC), or a nurse or other staff member trained in lactation support. Trained staff members may be able to perform much of the routinely provided breastfeeding support under the guidance of the pediatrician or other health care provider. The staff should be aware of community resources, including IBCLCs and other lactation support personnel, especially if one is not available in the practice (see recommendation 15).
3. Become knowledgeable regarding the rare but true contraindications to breastfeeding, which include infants with the classic form of galactosemia, maternal HIV or antiretroviral therapy, untreated active tuberculosis, human T-cell lymphotropic virus type I or II, use of illicit drugs, or mothers undergoing chemotherapy or radiation treatment.<sup>34</sup> Most maternal medications are compatible with breastfeeding. Specific drug information can be verified through the National Institutes of Health Toxicology

**TABLE 3** Summary of Breastfeeding Supportive Office Practices

- 
1. Have a written breastfeeding-friendly office policy
  2. Train staff in breastfeeding support skills
  3. Discuss breastfeeding during prenatal visits and at each well-child visit
  4. Encourage exclusive breastfeeding for ~6 months
  5. Provide appropriate anticipatory guidance that supports the continuation of breastfeeding as long as desired
  6. Incorporate breastfeeding observation into routine care
  7. Educate mothers on breast-milk expression and return to work
  8. Provide noncommercial breastfeeding educational resources for parents
  9. Encourage breastfeeding in the waiting room, but provide private space on request
  10. Eliminate the distribution of free formula
  11. Train staff to follow telephone triage protocols to address breastfeeding concerns
  12. Collaborate with the local hospital or birthing center and obstetric community regarding breastfeeding-friendly care
  13. Link with breastfeeding community resources
  14. Monitor breastfeeding rates in your practice
- 

Data Network, LactMed, which is accessible online or through a mobile device application.<sup>35</sup>

4. Introduce the subject of breastfeeding as early as possible, ideally with prenatal visits and early postpartum visits.<sup>27</sup> Use open-ended questions to inquire about feeding plans for the child, such as “What are your thoughts about feeding your baby?” or “Tell me about your previous infant feeding experiences.” Encourage attendance by both parents and/or partners at all visits, and consider discussions with grandparents or other important decision-makers in the family.<sup>36</sup>
5. Encourage breastfeeding mothers to feed newborn infants only human milk and to avoid offering supplements, including formula, glucose water, or other liquids, unless medically indicated.<sup>1</sup> This education ideally should begin prenatally, in anticipation of the newborn infant’s stay in the maternity hospital, and should continue through the early postnatal visits.
6. Work with committees within the local hospital or birthing center to implement breastfeeding-friendly care.<sup>1,17,18</sup>

Provide the hospital or birthing center with your office policies regarding breastfeeding. If the facility is not aware of the WHO/United Nations Children’s Fund Ten Steps to Successful Breastfeeding, the pediatrician can provide education and help to develop breastfeeding-friendly order sets for the hospital. Encourage the facility to pursue the Baby-Friendly USA designation so that mothers and infants are exposed to maternal and newborn care that supports and encourages breastfeeding, beginning with skin-to-skin care in the immediate postpartum period and early initiation of breastfeeding.<sup>20</sup> Show support for breastfeeding during hospital rounds by reinforcing the benefits of breastfeeding, encouraging exclusive breastfeeding, educating about the importance of frequent breastfeeding, and assessing the adequacy of breastfeeding. During rounds, either evaluate a feeding directly or review the chart for documentation of adequacy of feeding. Encourage mothers to attend breastfeeding classes. Advocate for lactation consultation for mothers who are experiencing any breastfeeding problems or who have concerns.



7. Schedule the first newborn visit by the third to fifth day of life, or approximately 24 to 48 hours from the time the newborn infant is discharged depending on the length of the hospital stay. At office visits, incorporate anticipatory guidance that supports exclusive breastfeeding until infants are approximately 6 months old, followed by continued breastfeeding for 1 year or longer, as mutually desired by the infant and mother.<sup>1</sup> Anticipatory guidance should include appropriate guidance about weight-gain expectations with the use of appropriate growth charts, such as the WHO growth standards for ages 0 to 2 years recommended by the CDC.<sup>37</sup>
8. Educate mothers regarding the provisions of the Patient Protection and Affordable Care Act (Pub L No. 111-148 [2010]), which cover access to breastfeeding support services, breaks to breastfeed or pump at work, as applicable,<sup>38</sup> and the ability to obtain breast pumps through insurance.<sup>39</sup>
9. Provide mothers with anticipatory guidance about returning to work. Workplace support in the pediatric practice and in other work environments can be optimized through the implementation of guidance from the Health Resources and Services Administration Maternal and Child Health Bureau's Business Case for Breastfeeding.<sup>32</sup> Provide information and education to mothers about both the expression and the storage of human milk, which may include providing parents with handouts detailing the recommendations regarding expression and storage.<sup>40</sup>
10. Have the front office staff advise the family, when the first follow-up appointment is scheduled, that the pediatrician or other trained office staff may wish to observe a feeding during the first visit, so that the family will be aware. Encourage the family to let the staff know when the infant is ready to feed while waiting for the appointment. For the first and subsequent appointments, feedings should be observed when the mother identifies any breastfeeding problem, or if weight gain is not appropriate.
11. Provide appropriate educational resources for parents.<sup>41,42</sup> These resources could cover, at a minimum, the benefits of breastfeeding for mother and child, AAP recommendations for duration of breastfeeding, education regarding feeding cues, how to tell whether the infant is getting enough milk, latch and holding techniques, and a list of peer support groups and local breastfeeding resources. The literature should be culturally sensitive and appropriate for the literacy of the patient population. Consider linking to appropriate resources on the practice Web site. Avoid distributing literature provided by manufacturers of infant formula.
12. Allow and encourage breastfeeding in the waiting room.<sup>33</sup> Display noncommercial posters and pamphlets that encourage mothers to breastfeed in waiting areas and examination rooms. Include graphics that show diversity and include fathers, who are valuable partners in the success of the breastfeeding dyad. Do not interrupt or discourage breastfeeding, either in the waiting room or in the examination room. Provide a comfortable, private area for mothers to breastfeed if they prefer privacy. This room may include a rocking chair, pillows, music, water fountain, or whatever helps to create a warm and relaxing environment. An examination room may suffice as a private room for breastfeeding.
13. Eliminate the practice of distribution of free formula and other infant items from formula companies to parents.<sup>43,44</sup> In accordance with the WHO International Code of Marketing of Breast-milk Substitutes,<sup>45</sup> the storage of formula supplies, which may be purchased by the practice as applicable for formula-fed infants, should be out of the view of patients. The breastfeeding-friendly pediatric office practice should not accept gifts (formula and other feeding supplies, pens, writing pads, calendars, mugs, etc) from companies manufacturing infant formula, feeding bottles, or pacifiers. Consumer publications that advertise infant formula or have tear-off cards or inserts to receive free or discounted formula should be discouraged.
14. Train staff to follow telephone triage protocols to address breastfeeding concerns and problems. Train staff on providing appropriate breastfeeding telephone advice, including when to refer to an IBCLC or to a physician with special expertise in breastfeeding management.<sup>46</sup> Telehealth consults may be an option in some locations.
15. Acquire or maintain a list of community resources and be knowledgeable about referral procedures. Refer expectant and new parents to peer, community support, and resource groups.<sup>33</sup> Get to know peer and community support groups in your area. WIC breastfeeding support services, La Leche League International, and peer counselors are

some options. Not all groups may provide evidence-based information, so it is useful if the pediatric care provider or members of the clinic staff have attended and maintain good communication with the leaders of these support groups. Consider inviting successfully breastfeeding mothers to join any advisory groups associated with the practice.

16. Frenulum clipping of the tongue/upper lip has become a popular practice, which may improve infant latch and the effectiveness of breastfeeding and milk transfer. Work closely with local lactation consultants or breastfeeding specialists to determine if frenotomy is appropriate. If so, refer to otolaryngologists and/or dentists with experience in clipping the frenulum in a timely fashion, if the physicians in the practice do not perform the procedure.
17. Collaborate with the obstetric community to develop optimal breastfeeding support programs, because it is clearly documented that a mother's decision to breastfeed starts in the prenatal period and, in many cases, before pregnancy.<sup>10,36</sup>
18. Provide support and education to local child care centers on the importance of breastfeeding and the handling, storage, and feeding of expressed human milk.<sup>47</sup>
19. Monitor breastfeeding initiation and duration rates in the pediatric practice. Be able to access state and national trends as tracked by the CDC through the National Immunization Survey<sup>11</sup> and hospital practices through the Maternity Practices in Infant Nutrition and Care biannual survey data.<sup>48</sup>

## **BARRIERS TO IMPLEMENTING BREASTFEEDING-FRIENDLY OFFICE PRACTICES**

There are challenges to implementing the steps as outlined above for the breastfeeding-friendly pediatric office practice. Breastfeeding care delivered by physicians in the outpatient practice may be limited by lack of knowledge, skills, time, and cultural sensitivity.<sup>23</sup> Breastfeeding management and counseling can be both time consuming and labor intensive. In many health care systems, especially in the United States, practice revenue currently is dependent on volume of visits and patients seen instead of the outcomes of those patients. The AAP Section on Breastfeeding has developed a guide on coding and billing<sup>49</sup> as a tool to optimize reimbursement for time spent in breastfeeding support in the office. Coding guidance also is available through the American College of Obstetricians and Gynecologists for maternal conditions.<sup>50</sup> If additional office visits are required beyond those covered for well-child visits, then diagnosis codes should be used for conditions such as jaundice, newborn feeding problem, infection associated with lactation, etc. Extended visits that require repeated evaluation of breastfeeding and extensive counseling should be billed according to time-based codes.

Even if payment were not an issue, time constraints may be. Complicated breastfeeding problems may require immediate attention and can monopolize staff time and space. Some of these more complex issues may exceed the knowledge level or skill set of the general pediatric care provider. Identifying health care providers and lactation support personnel in the community as sources for referral is important for timely intervention. Lactation support personnel may be limited in certain geographic distributions, especially

rural areas, as well as in certain communities. There are different skill levels among lactation support personnel. These may include, among others, volunteer or paid peer counselors, lactation educators, certified lactation counselors, IBCLCs, and breastfeeding medicine specialists. The extent of training required by the lactation specialist may vary depending on the needs of the mother and/or infant and the particular clinical or community setting. There exists a need for greater availability of culturally appropriate and ethnically diverse lactation support personnel in a variety of community and health care settings. Home visitation, where available, is a family-friendly way to provide lactation support. Pediatric care providers need to be knowledgeable about common complications that can affect both mother and infant and either treat these conditions in the mother or become aware of obstetrician/gynecologists, family physicians, or breastfeeding medicine specialists in the community who are comfortable and knowledgeable in treating the maternal conditions.

Clearly, there will be obstacles to implementing the steps outlined in this report; however, these obstacles are not insurmountable. With ongoing advocacy for the support of breastfeeding, there should be fewer barriers over time.

## **FURTHER RESEARCH**

It would be valuable to study the cost-effectiveness and impact of the routine presence of trained lactation support personnel, such as IBCLCs, versus no specific lactation support in the outpatient pediatric practice on breastfeeding exclusivity and duration. In addition, further research is needed on the effectiveness and intensity of other interventions to support breastfeeding, such as in-person

consultation, group sessions, online resources, or use of telehealth, as well as community support interventions. Another area for research is the best method for improving pediatric care provider education in breastfeeding management and support during undergraduate and graduate medical education, as well as continuing medical education for practicing physicians. Education of other members of the health care team who may interact with breastfeeding mothers or children is also important. Finally, similar to the research published in *Pediatrics* entitled “Evaluation of an Office Protocol To Increase Exclusivity of Breastfeeding,”<sup>28</sup> in which the authors evaluated the ABM’s clinical protocol for the Breastfeeding-Friendly Physician’s Office,<sup>27</sup> an evaluation of the impact of implementation in pediatric office practices of the steps outlined in this clinical report on rates of exclusive breastfeeding and duration of breastfeeding would be beneficial.

## CONCLUSIONS

The benefits of breastfeeding and the potential risks of not breastfeeding are numerous, and increasing

breastfeeding initiation, duration, and exclusivity has been the focus of multiple recent initiatives in the health care setting, workplace, and community, as recommended by the *Surgeon General’s Call to Action To Support Breastfeeding*. With national goals as outlined in Healthy People 2020, an increase in overall breastfeeding initiation, and more Baby-Friendly–designated hospitals, the need for increased support from all members of the health care team is clear. The pediatric care provider is well suited to play a primary role in this effort. The steps outlined in this document provide clear and concise ways for the pediatric office practice to support breastfeeding mothers, infants, and families; increase breastfeeding exclusivity and duration in their patients; and improve health outcomes for the population.

## AUTHORS

Joan Younger Meek, MD, MS, RD, FAAP, IBCLC  
Amy J. Hatcher, MD, FAAP

## SECTION ON BREASTFEEDING EXECUTIVE COMMITTEE, 2015–2016

Joan Younger Meek, MD, MS, RD, FAAP, IBCLC,  
Chairperson

Margreete Johnston, MD, MPH, FAAP  
Mary O’Connor, MD, MPH, FAAP  
Lisa Stellwagen, MD, FAAP  
Jennifer Thomas, MD, MPH, FAAP  
Julie Ware, MD, FAAP  
Richard Schanler, MD, FAAP, Immediate Past  
Chairperson

## SUBCOMMITTEE CHAIRPERSONS

Lawrence Noble, MD, FAAP – Policy Chairperson  
Krystal Revai, MD, MPH, FAAP – Chief Chapter  
Breastfeeding Coordinator  
Natasha Sriraman, MD, MPH, FAAP – Education  
Program Chairperson

## STAFF

Ngozi Onyema-Melton, MPH, CHES

## ABBREVIATIONS

AAP: American Academy of  
Pediatrics  
ABM: Academy of Breastfeeding  
Medicine  
CDC: Centers for Disease Control  
and Prevention  
IBCLC: International Board  
Certified Lactation  
Consultant  
WHO: World Health Organization  
WIC: Special Supplemental  
Nutrition Program for  
Women, Infants, and  
Children

**FUNDING:** No external funding.

**POTENTIAL CONFLICT OF INTEREST:** The authors have indicated they have no potential conflicts of interest to disclose.

## REFERENCES

1. Eidelman AI, Schanler RJ, Johnston M, et al; Section on Breastfeeding. Breastfeeding and the use of human milk. *Pediatrics*. 2012;129(3). Available at: [www.pediatrics.org/cgi/content/full/129/3/e827](http://www.pediatrics.org/cgi/content/full/129/3/e827)
2. Chen A, Rogan WJ. Breastfeeding and the risk of postneonatal death in the United States. *Pediatrics*. 2004;113(5). Available at: [www.pediatrics.org/cgi/content/full/113/5/e435](http://www.pediatrics.org/cgi/content/full/113/5/e435)
3. Grummer-Strawn LM, Rollins N. Summarising the health effects of breastfeeding. *Acta Paediatr*. 2015;104(467):1–2
4. Collaborative Group on Hormonal Factors in Breast Cancer. Breast cancer and breastfeeding: collaborative reanalysis of individual data from 47 epidemiological studies in 30 countries, including 50302 women with breast cancer and 96973 women without the disease. *Lancet*. 2002;360(9328):187–195
5. Stuebe AM, Rich-Edwards JW, Willett WC, Manson JE, Michels KB. Duration of lactation and incidence of type 2 diabetes. *JAMA*. 2005;294(20):2601–2610
6. Schwarz EB, Ray RM, Stuebe AM, et al. Duration of lactation and risk factors for maternal cardiovascular disease. *Obstet Gynecol*. 2009;113(5):974–982
7. Victora CG, Bahl R, Barros AJ, et al; Lancet Breastfeeding Series Group. Breastfeeding in the 21st century: epidemiology, mechanisms, and lifelong effect. *Lancet*. 2016;387(10017):475–490
8. US Department of Health and Human Services. The Surgeon General’s Call to Action To Support Breastfeeding. Washington, DC: US Department of Health and Human Services, Office of the Surgeon General; 2011. Available at: [www.surgeongeneral.gov/library/](http://www.surgeongeneral.gov/library/)

- calls/breastfeeding/. Accessed August 10, 2016
9. Academy of Breastfeeding Medicine Board of Directors. Position on breastfeeding. *Breastfeed Med.* 2008;3(4):267–270
  10. American College of Obstetricians and Gynecologists. Optimizing support for breastfeeding as part of obstetric practice. Committee Opinion No. 658. *Obstet Gynecol.* 2016;127(2):e86–e92
  11. American College of Obstetricians and Gynecologists. Breastfeeding in underserved women: increasing initiation and continuation of breastfeeding. Committee Opinion No. 570. *Obstet Gynecol.* 2013; 122(2 pt 1):423–428
  12. American Academy of Family Physicians. Breastfeeding, family physicians supporting [position paper]. Available at: [www.aafp.org/about/policies/all/breastfeeding-support.html](http://www.aafp.org/about/policies/all/breastfeeding-support.html). Accessed August 10, 2016
  13. Centers for Disease Control and Prevention. National immunization survey. Available at: [www.cdc.gov/breastfeeding/data/nis\\_data/rates-any-exclusive-bf-socio-dem-2013.htm](http://www.cdc.gov/breastfeeding/data/nis_data/rates-any-exclusive-bf-socio-dem-2013.htm). Accessed August 10, 2016
  14. US Department of Health and Human Services. Healthy People 2020: maternal, infant, and child health. Available at: <https://www.healthypeople.gov/2020/topics-objectives/topic/maternal-infant-and-child-health>. Accessed August 10, 2016
  15. Perrine CG, Scanlon KS, Li R, Odom E, Grummer-Strawn LM. Baby-friendly hospital practices and meeting exclusive breastfeeding intention. *Pediatrics.* 2012;130(1):54–60
  16. The Joint Commission. Perinatal care core measures. Available at: [www.jointcommission.org/perinatal\\_care/](http://www.jointcommission.org/perinatal_care/). Accessed August 10, 2016
  17. National Institute for Children’s Health Quality. Projects: Best Fed Beginnings. Available at: <http://breastfeeding.nichq.org/solutions/best-fed-beginnings>. Accessed August 10, 2016
  18. World Health Organization. Evidence for the Ten Steps to Successful Breastfeeding. Available at: [www.who.int/maternal\\_child\\_adolescent/documents/9241591544/en/](http://www.who.int/maternal_child_adolescent/documents/9241591544/en/). Accessed August 10, 2016
  19. World Health Organization. Baby-Friendly Hospital Initiative: revised, updated and expanded for integrated care. Available at: [www.who.int/nutrition/publications/infantfeeding/bfhi\\_trainingcourse/en/](http://www.who.int/nutrition/publications/infantfeeding/bfhi_trainingcourse/en/). Accessed August 10, 2016
  20. World Health Organization. Baby-Friendly Hospital Initiative. Available at: [www.who.int/nutrition/topics/bfhi/en/](http://www.who.int/nutrition/topics/bfhi/en/). Accessed August 10, 2016
  21. EMPower Breastfeeding [home page]. Available at: <http://empowerbreastfeeding.org/>. Accessed August 10, 2016
  22. Baby-Friendly USA [home page]. Available at: <https://www.babyfriendlyusa.org/>. Accessed August 10, 2016
  23. Feldman-Winter LB, Schanler RJ, O’Connor KG, Lawrence RA. Pediatricians and the promotion and support of breastfeeding. *Arch Pediatr Adolesc Med.* 2008;162(12):1142–1149
  24. Szucs KA, Miracle DJ, Rosenman MB. Breastfeeding knowledge, attitudes, and practices among providers in a medical home. *Breastfeed Med.* 2009;4(1):31–42
  25. Academy of Breastfeeding Medicine. Educational objectives and skills for the physician with respect to breastfeeding. *Breastfeed Med.* 2011;6(2):99–105
  26. US Breastfeeding Committee. *Core Competencies in Breastfeeding Care and Services for All Health Professionals, Revised Edition.* Washington, DC: US Breastfeeding Committee; 2010
  27. Grawey AE, Marinelli KA, Holmes AV; Academy of Breastfeeding Medicine. ABM clinical protocol #14. Breastfeeding-friendly physician’s office: optimizing care for infants and children, revised 2013. *Breastfeed Med.* 2013;8(2):237–242
  28. Corriveau SK, Drake EE, Kellams AL, Rovnyak VG. Evaluation of an office protocol to increase exclusivity of breastfeeding. *Pediatrics.* 2013;131(5):942–950
  29. Feldman-Winter L. Evidence-based interventions to support breastfeeding. *Pediatr Clin North Am.* 2013;60(1):169–187
  30. Chung M, Raman G, Trikalinos T, Lau J, Ip S. Interventions in primary care to promote breastfeeding: an evidence review for the U.S. Preventive Services Task Force. *Ann Intern Med.* 2008;149(8):565–582
  31. US Preventive Services Task Force. Final recommendation statement: breastfeeding: counseling. Available at: [www.uspreventiveservicestaskforce.org/Page/Document/RecommendationStatementFinal/breastfeeding-counseling](http://www.uspreventiveservicestaskforce.org/Page/Document/RecommendationStatementFinal/breastfeeding-counseling). Accessed August 10, 2016
  32. US Department of Health and Human Services; Health Resources and Service Administration, Maternal and Child Health Bureau. The business case for breastfeeding. Available at: [www.womenshealth.gov/breastfeeding/government-in-action/business-case-for-breastfeeding/business-case-for-breastfeeding-for-business-managers.pdf](http://www.womenshealth.gov/breastfeeding/government-in-action/business-case-for-breastfeeding/business-case-for-breastfeeding-for-business-managers.pdf). Accessed August 10, 2016
  33. American Academy of Pediatrics Section on Breastfeeding. 10 Steps to support parent’s choice to breastfeed their baby. Available at: <http://www2.aap.org/breastfeeding/files/pdf/TenStepsPoster.pdf>. Accessed August 10, 2016
  34. Centers for Disease Control and Prevention. Breastfeeding: diseases and conditions. When should a mother avoid breastfeeding? Available at: [www.cdc.gov/breastfeeding/disease/index.htm](http://www.cdc.gov/breastfeeding/disease/index.htm). Accessed August 10, 2016
  35. National Institutes of Health; US National Library of Medicine. LactMed: a TOXNET database. Available at: <http://toxnet.nlm.nih.gov/newtoxnet/lactmed.htm>. Accessed August 10, 2016
  36. American College of Obstetricians and Gynecologists. Physician conversation guide on support for breastfeeding. Available at: [www.acog.org/About-ACOG/ACOG-Departments/Toolkits-for-Health-Care-Providers/Breastfeeding-Toolkit/Physician-Conversation-Guide-on-Support-for-Breastfeeding](http://www.acog.org/About-ACOG/ACOG-Departments/Toolkits-for-Health-Care-Providers/Breastfeeding-Toolkit/Physician-Conversation-Guide-on-Support-for-Breastfeeding). Accessed August 10, 2016
  37. Centers for Disease Control and Prevention. The WHO growth charts. Available at: [www.cdc.gov](http://www.cdc.gov).



- gov/growthcharts/who\_charts.htm#The%20WHO%20Growth%20Charts. Accessed August 10, 2016
38. US Department of Labor, Wage and Hour Division. Break time for nursing mothers. Available at: [www.dol.gov/whd/nursingmothers/](http://www.dol.gov/whd/nursingmothers/). Accessed August 10, 2016
  39. HealthCare.gov. Breastfeeding benefits. Available at: <https://www.healthcare.gov/coverage/breast-feeding-benefits/>. Accessed August 10, 2016
  40. Academy of Breastfeeding Medicine Protocol Committee. ABM clinical protocol #8: human milk storage information for home use for full-term infants. *Breastfeed Med*. 2010;5(3):127–130
  41. US Health and Human Services Office on Women's Health. Breastfeeding. Available at: [www.womenshealth.gov/breastfeeding/](http://www.womenshealth.gov/breastfeeding/). Accessed August 10, 2016
  42. Healthychildren.org; American Academy of Pediatrics. Breastfeeding. Available at: [ages-stages/baby/breastfeeding/](https://www.healthychildren.org/English/ages-stages/baby/breastfeeding/). Accessed August 10, 2016
  43. Rosenberg KD, Eastham CA, Kasehagen LJ, Sandoval AP. Marketing infant formula through hospitals: the impact of commercial hospital discharge packs on breastfeeding. *Am J Public Health*. 2008;98(2):290–295
  44. Howard C, Howard F, Lawrence R, Andresen E, DeBlieck E, Weitzman M. Office prenatal formula advertising and its effect on breastfeeding patterns. *Obstet Gynecol*. 2000;95(2):296–303
  45. World Health Organization. International code of marketing of breast-milk substitutes. Available at: [www.unicef.org/nutrition/files/nutrition\\_code\\_english.pdf](http://www.unicef.org/nutrition/files/nutrition_code_english.pdf). Accessed August 10, 2016
  46. Bunik M. *Breastfeeding Telephone Triage and Advice*. 2nd ed. Elk Grove Village, IL: American Academy of Pediatrics; 2016
  47. UNC Gillings School of Global Public Health; Carolina Global Breastfeeding Institute. Tools for action-breastfeeding and child care. Available at: <http://breastfeeding.sph.unc.edu/what-we-do/programs-and-initiatives/breastfeeding-and-child-care-toolkit/>. Accessed August 10, 2016
  48. Centers for Disease Control and Prevention. Maternity Practices in Infant Nutrition and Care (mPINC) survey. Available at: [www.cdc.gov/breastfeeding/data/mpinc/index.htm](http://www.cdc.gov/breastfeeding/data/mpinc/index.htm). Accessed August 10, 2016
  49. American Academy of Pediatrics. Supporting breastfeeding and lactation: the primary care pediatrician's guide to getting paid. Available at: <http://www2.aap.org/breastfeeding/files/pdf/coding.pdf>. Accessed August 10, 2016
  50. American College of Obstetricians and Gynecologists. Breastfeeding coding. Available at: [www.acog.org/About-ACOG/ACOG-Departments/Toolkits-for-Health-Care-Providers/Breastfeeding-Toolkit/Breastfeeding-Coding](http://www.acog.org/About-ACOG/ACOG-Departments/Toolkits-for-Health-Care-Providers/Breastfeeding-Toolkit/Breastfeeding-Coding). Accessed August 10, 2016

**The Breastfeeding-Friendly Pediatric Office Practice**  
Joan Younger Meek, Amy J. Hatcher and SECTION ON BREASTFEEDING  
*Pediatrics*; originally published online April 17, 2017;  
DOI: 10.1542/peds.2017-0647

<b>Updated Information &amp; Services</b>	including high resolution figures, can be found at: <a href="/content/early/2017/04/13/peds.2017-0647.full.html">/content/early/2017/04/13/peds.2017-0647.full.html</a>
<b>References</b>	This article cites 21 articles, 3 of which can be accessed free at: <a href="/content/early/2017/04/13/peds.2017-0647.full.html#ref-list-1">/content/early/2017/04/13/peds.2017-0647.full.html#ref-list-1</a>
<b>Subspecialty Collections</b>	This article, along with others on similar topics, appears in the following collection(s): <b>Administration/Practice Management</b> <a href="/cgi/collection/administration:practice_management_sub">/cgi/collection/administration:practice_management_sub</a> <b>Nutrition</b> <a href="/cgi/collection/nutrition_sub">/cgi/collection/nutrition_sub</a> <b>Breastfeeding</b> <a href="/cgi/collection/breastfeeding_sub">/cgi/collection/breastfeeding_sub</a>
<b>Permissions &amp; Licensing</b>	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: <a href="/site/misc/Permissions.xhtml">/site/misc/Permissions.xhtml</a>
<b>Reprints</b>	Information about ordering reprints can be found online: <a href="/site/misc/reprints.xhtml">/site/misc/reprints.xhtml</a>

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2017 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



# PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

## **The Breastfeeding-Friendly Pediatric Office Practice**

Joan Younger Meek, Amy J. Hatcher and SECTION ON BREASTFEEDING

*Pediatrics*; originally published online April 17, 2017;

DOI: 10.1542/peds.2017-0647

The online version of this article, along with updated information and services, is located on the World Wide Web at:

[/content/early/2017/04/13/peds.2017-0647.full.html](http://content.early/2017/04/13/peds.2017-0647.full.html)

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2017 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

